This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.




Figure 1: General structure of activators of non-genomic Estrogen-Like Signalling (ANGELS)





Figure 2: Estrogen deficiency causes increased apoptosis of osteoblasts and osteocytes in murine vertebral bone.



::

Figure 3: Inhibition of apoptosis of osteoblastic cells.




Figure 4: Inhibition of apoptosis of MLO-Y4 osteocytic cells by ANGELS





Figure 5: Blockade of the anti-apoptotic effect of estrogen and ANGELS by ICI 182,780 in osteoblastic cells






Figure 7: Estrogen receptor a or b is required for the antiapoptotic effects of 17b estradiol, 17a estradiol, and estratriene-3-of on etoposide-induced apoptosis (experiment 1/21//99).




Figure 8: Activation of Extracellular Signal Regulated Kinases (ERKs)




Figure 9: The effect of estrogenic compounds on the activation of ERK1/2 is blocked by specific inhibitor.




Figure 10: The specific inhibitor of ERK activation abolishes the anti-apoptotic effect of the estrogenic compounds.






Figure 11 : Unlike 17 β estradiol, estratriene-3-ol does not transactivate an estrogen response element through ERlpha.



[2S-(2a,4aα,10aβ)]-1,2,3,4,4a,9,10,10a-Octahydro-7-hydroxy-2-methyl-2-phenanthrenemethanol

[2S-(2a,4aα,10aβ)]-1,2,3,4,4a,9,10,10a-Octahydro-7-hydroxy-2-methyl-2-phenanthrenecarboxaldehyde

 $C_{16}H_{20}O_2$ MW=244







Figure 14: Mechanisms of Estrogen Receptor Action

Formation occurs only on sites of previous osteoclastic bone resorption.



Anti-fracture efficacy (through inhibition of osteocyte apoptosis)

Figure 15: Implications of the effects of anti-resorptive vs. non anti-resorptive agents on apoptosis

	R AND/OR h	SUBSTITUTION.						
		STRUCTURE						
- HYCT.		-O l						
) ree		-CH ₃						
METHYL ET		-0CH3						
ACE		0-C-CH3						
EIML ET		0-CH2-CH3						
3. 3. (OF 17		OCH3						
DIMETHYL FL		OCH ₃						
हामा≁		_C ≡ CH						
		\Q						
BENT		کسر '۔ن						
BENZYL ET	, , , , , , , , , , , , , , , , , , , ,	00H2(D)						
GLÚCURO:		C ₆ H ₈ O ₆						
SULFATE SODIUM :		0503Nc						
C:::		=						
VALE		-C5HgC						
CYCLOPENTYLPROPIE		-0-0-(CH ₂) ₂ -						
PROPIN		-0-C-(CH ₂) ₂						
HEMISUCC"		-C4 H4 O2						
PALA:		-C16H32O2						

Figure 16A

Ř.	AND/OR	Ro	SUBSTITUTIONS				
			STRUCTURE				
SODIUM PHOSPI			-0-P03No2				
ENANTH!			-C7H120				
GLUCURONIDE SODIUM	S	-C6H8O6Nc					
STEAT			-C18H340				
TRETHYL" AMMONIUM S		-N-(C2H5)3					
CYPIC			0-C-CH2CH2 -				

Figure 16B

Name 1	3,1 P-Estradol	3,17REsandal 3-O-ME	डिएक्ट्रीकारु ३-०।	3,17c.Esman	3,178-Benufol 3-scence	2-ffydoxy-17\$-ecuafol	17 Berensol 2.3-CME	Estrace	Estrone 3-O-ME	ट् स्मेल	Estriol 3-O-ME		Ethynyl Estradol	Metracol
	R	R=CH3	R=A	R	R=CEsCO	## #	R=CH3	R=H	RaCH ₃	Ref	R=CH5		H-R-H	R=CH3
	***			E R	\	No. 2		\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	}			P. O.	A POLICE CH	




Figure 19: Effect of the all peptide on the 17bE₂-induced ERE activity in 293 cells




Figure 20: Effect of the all peptide on the 17bE₂-induced inhibition of IL-6 activity in 293 cells



